

PART B — (5 × 16 = 80 marks)

11. (a) (i) Find the eigenvalues and eigenvectors of the matrix $\begin{pmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{pmatrix}$. (8)

- (ii) Using Cayley–Hamilton theorem find A^{-1} , where $A = \begin{pmatrix} 7 & 2 & -2 \\ -6 & -1 & 2 \\ 6 & 2 & -1 \end{pmatrix}$. (8)

Or

- (b) Reduce the quadratic form $6x_1^2 + 3x_2^2 + 3x_3^2 - 4x_1x_2 - 2x_2x_3 + 4x_3x_1$ to canonical form. (16)

12. (a) (i) Examine the convergence of the series $\frac{1}{2!} - \frac{2}{3!} + \frac{3}{4!} \dots \infty$. (8)

- (ii) Find the sum to infinity of the series $\frac{1}{1!} + \frac{1+5}{2!} + \frac{1+5+5^2}{3!} + \dots \infty$. (8)

Or

- (b) (i) Expand $\frac{1}{(1-2x)^2(1-3x)}$ in ascending powers of x . Also find the coefficient of x^2 . (8)

- (ii) Prove that $\sqrt{x^2+4} - \sqrt{x^2+1} = 1 - \frac{x^2}{4} + \frac{7}{64}x^4$ nearly when x is small. (8)

13. (a) (i) Find the equation of the circle of curvature of the parabola $y^2 = 12x$ at $(3, 6)$. (8)

- (ii) Find the equation of evolute of the curve $x = a(\cos t + t \sin t)$, $y = a(\sin t - t \cos t)$. (8)

Or

- (b) (i) Find the radius of curvature at $(a, 0)$ on the curve $xy^2 = a^3 - x^3$. (8)

- (ii) Find the envelope of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where a and b are connected by the relation $a^2 + b^2 = c^2$, c being a constant. (8)

14. (a) (i) If $u = f(r, s, t)$ and $r = \frac{x}{y}, s = \frac{y}{z}, t = \frac{z}{x}$, find the value of $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z}$. (8)

(ii) Examine the extrema of $f(x, y) = x^2 + xy + y^2 + \frac{1}{x} + \frac{1}{y}$. (8)

Or

(b) (i) Using Taylor's series expansion, expand $e^x \sin y$ in powers of x and y as far as terms of the 3rd degree. (8)

(ii) Find the shortest and longest distances from the point $(1, 2, -1)$ to the sphere $x^2 + y^2 + z^2 = 24$. (8)

15. (a) (i) Evaluate $\int_0^{a\sqrt{a^2-x^2}} \int_0^{\sqrt{a^2-x^2-y^2}} \sqrt{a^2-x^2-y^2} \, dx \, dy$. (8)

(ii) Using double integral find the area of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. (8)

Or

(b) (i) Change the order of integration in $\int_0^{2\sqrt{4-y^2}} \int_0^y xy \, dx \, dy$ and evaluate it. (8)

(ii) By transforming into polar co-ordinates evaluate $\iint_{00}^{\infty\infty} e^{-(x^2+y^2)} \, dx \, dy$.

Hence find the value of $\int_0^{\infty} e^{-x^2} \, dx$. (8)