

Reg. No. :

Question Paper Code : 90821

B.E./B.Tech. DEGREE EXAMINATIONS, NOVEMBER/DECEMBER 2022.

Fourth/Fifth/Sixth Semester

Aeronautical Engineering

MA 8491 – NUMERICAL METHODS

(Common to : Aerospace Engineering/Agriculture Engineering/Civil Engineering/Electrical and Electronics Engineering/Electronics and Instrumentation Engineering/Instrumentation and Control Engineering/Manufacturing Engineering/Mechanical Engineering (Sandwich)/Mechanical and Automation Engineering/Biotechnology and Biochemical Engineering/Chemical Engineering/Chemical and Electrochemical Engineering/Plastic Technology/Polymer Technology/Textile technology)

(Regulations 2017)

Time : Three hours

Maximum : 100 marks

Answer ALL questions.

PART A — (10 × 2 = 20 marks)

1. State the sufficient condition for the convergence of Newton-Raphson method for the equation $f(x) = 0$.
2. State the principle used in Gauss Jordan method.
3. Find the second divided difference with arguments a , b and c of the function $f(x) = \frac{1}{x}$.
4. What are the advantages of Lagrange's formula over Newton's forward and backward interpolation formulae?
5. Write the formula for $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ at $x = x_n$ by using Newton's backward difference operator.


6. What is the restriction on the number of intervals in order to evaluate $\int_a^b f(x) dx$ by Trapezoidal rule and by Simpson's one-third rule?
7. State the modified Euler formula to find $y(x_1)$ for solving $\frac{dy}{dx} = f(x, y)$, $y(x_0) = y_0$.
8. Write down Adam-Bashforth predictor and corrector formulae.
9. Write down the finite difference scheme for solving $\frac{d^2y}{dx^2} = x + y$, $y(0) = 0 = y(1)$ with $h = 0.5$.
10. State the explicit formula for the one dimensional wave equation $u_{tt} = \alpha^2 u_{xx}$ with $1 - \lambda^2 \alpha^2 = 0$, where $\lambda = \frac{k}{h}$.

PART B — (5 × 16 = 80 marks)

11. (a) (i) Find the smallest positive root of $x^3 - 2x - 5 = 0$ by the fixed point iteration method, correct to three decimal places. (6)
- (ii) Find all eigenvalues and the corresponding eigenvectors of a matrix $A = \begin{pmatrix} 1 & \sqrt{2} & 2 \\ \sqrt{2} & 3 & \sqrt{2} \\ 2 & \sqrt{2} & 1 \end{pmatrix}$ by Jacobi's method. (10)

Or

- (b) (i) Find, by Power method, the largest eigenvalue and the corresponding eigenvector of a matrix $A = \begin{pmatrix} 1 & 6 & 1 \\ 1 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}$ starting with initial vector $X^{(0)} = (1 \ 0 \ 0)^T$. (8)
- (ii) Solve the following system of equations by Gauss-Seidel method, correct to three decimal places :
 $28x + 4y - z = 32$; $x + 3y + 10z = 24$ and $2x + 17y + 4z = 35$. (8)
[perform 4 iterations in each above 4 questions]



(b) (i) Solve $\frac{dy}{dx} = \frac{y-x}{y+x}$, $y(0)=1$ by Euler's method to find $y(0.1)$ with $h = 0.05$. (6)

(ii) Solve $\frac{dy}{dx} = x - y^2$, $y(0) = 0$, $y(0.2) = 0.02$, $y(0.4) = 0.0795$ and $y(0.6) = 0.1762$ by Milne's method to find $y(0.8)$. (10)

15. (a) (i) Solve the Poisson equation $u_{xx} + u_{yy} = -81xy$, for $0 < x, y < 1$, given $u(x, 0) = 0 = u(0, y)$, and $u(x, 1) = 100 = u(1, y)$. (8)

(ii) Use Crank-Nicholson implicit scheme to solve $16 \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$, $0 < x < 1$ and $t > 0$ given $u(x, 0) = 0 = u(0, t)$, and $u(1, t) = 100t$. Compute $u(x, t)$ for one time step, taking $\Delta X = 0.25$. (8)

Or

(b) (i) Solve the boundary value problem $x y'' + y = 0$ with the boundary conditions $y(1)=1$ and $y(2)=2$, taking $h=1/4$ by finite difference method. (8)

(ii) Solve $u_{xx} + u_{yy} = 0$ for the following square mesh with boundary values as shown in the figure below (8)

